

References

1. DeFrances CJ, Cullen KA, Kozak LJ. National Discharge Survey: 2005 annual summary with detailed diagnosis and procedure data. National Center for Health Statistics. *Vital Health Stat 13*. 2007;(165):1-209.
2. Samelson EJ, Zhang Y, Kiel DP, et al. Effect of birth cohort on risk of hip fracture: age-specific incidence rates in the Framingham Study. *Am J Public Health*. 2002;92:858-862.
3. Zuckerman JD. Hip fracture. *N Engl J Med*. 1996;334:1519-1525.
4. Leibson CL, Tosteson ANA, Gabriel SE, et al. Mortality, disability, and nursing home use for persons with and without hip fracture: a population-based study. *J Am Geriatr Soc*. 2002;50:1644-1650.
5. Magaziner J, Hawkes W, Hebel JR, et al. Recovery from hip fracture in eight areas of function. *J Gerontol A Biol Sci Med Sci*. 2000;55:M498-M507.
6. Gdalevich M, Cohen D, Yosef D, Tauber C. Morbidity and mortality after hip fracture: the impact of operative delay. *Arch Orthop Trauma Surg*. 2004;124:334-340.
7. Phy MP, Vanness DJ, Melton LJ 3rd, et al. Effects of a hospital model on elderly patients with hip fracture. *Arch Intern Med*. 2005;165:796-801.
8. Batsis JA, Phy MP, Melton LJ 3rd, et al. Effects of a hospitalist care model on mortality of elderly patients with hip fractures. *J Hosp Med*. 2007;2:219-225.
9. Roy A, Heckman MG, Roy V. Associations between the hospitalist model of care and quality-of-care-related outcomes in patients undergoing hip fracture surgery. *Mayo Clin Proc*. 2006;81:28-31.
10. Huddleston JM, Long KH, Naessens JM, et al. Medical and surgical comanagement after elective hip and knee arthroplasty: a randomized, controlled trial. *Ann Intern Med*. 2004;141:28-38.
11. Canadian Orthopedic Foundation. Hip fractures. Available at: http://www.canorth.org/en/Images/ReferenceImages/HipInjuries_EN_sm.jpg. Accessed November 20, 2008.
12. Barsoum WK, Helfand R, Krebs V, Whinney C. Managing perioperative risk in the hip fracture patient. *Cleve Clin J Med*. 2006;73(suppl 1):S46-S50.

13. eOrthopod. Available at:
http://www.eorthopod.com/images/ContentImages/hip/hip_pinning/hip_pinning_procedure01.jpg. Accessed November 20, 2008.¹³
14. eOrthopod. Available at:
http://www.eorthopod.com/images/ContentImages/hip/hip_fracture/hip_fracture_treatment06.jpg. Accessed November 20, 2008.
15. Canadian Orthopedic Foundation. Hip fractures. Available at:
http://www.canorth.org/en/Images/ReferenceImages/HipRepairs_EN_sm.jpg. Accessed November 20, 2008.
16. Whinney CM. Do hip fractures need to be repaired within 24 hours of injury? *Cleve Clin J Med*. 2005;72:250-252.
17. eOrthopod. Available at:
http://www.eorthopod.com/images/ContentImages/hip/hip_fracture/hip_fracture_treatment01.jpg and
http://www.eorthopod.com/images/ContentImages/hip/hip_fracture/hip_fracture_treatment03.jpg. Accessed November 24, 2008.
18. Grimes JP, Gregory PM, Noveck H, et al. The effects of time-to-surgery on mortality and morbidity in patients following hip fracture. *Am J Med*. 2002;112:702-709.
19. Orosz GM, Magaziner J, Hannan EL, et al. Association of timing of surgery for hip fracture and patient outcomes. *JAMA*. 2004;291:1738-1743.
20. Al-Ani AN, Samuelsson B, Tidermark J, et al. Early operation on patients with a hip fracture improved the ability to return to independent living. *J Bone Joint Surg Am*. 2008;90:1436-1442.
21. Bottle A, Aylin P. Mortality associated with delay in operation after hip fracture: observational study. *BMJ*. 2006;332:947-951.
22. Morrison RS, Chassin MR, Siu AL. The medical consultant's role in caring for patients with hip fracture. *Ann Intern Med*. 1998;128:1010-1020.
23. Shiga T, Wajima Z, Ohe Y. Is operative delay associated with increased mortality of hip fracture patients? Systematic review, meta-analysis, and meta-regression. *Can J Anesth*. 2008;55:146-154.

24. Grant PJ, Wesorick DH. Perioperative medicine for the hospitalized patient. *Med Clin N Am.* 2008;92:325-348.
25. Lee TH, Marcantonio ER, Mangione CM, et al. Derivation and prospective validation of a simple index for prediction of cardiac risk of major noncardiac surgery. *Circulation.* 1999;100:1043-1049.
26. Fleisher LA, Beckman JA, Brown KA, et al. ACC/AHA 2007 guidelines on perioperative cardiovascular evaluation and care for noncardiac surgery: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 2002 Guidelines on Perioperative Cardiovascular Evaluation for Noncardiac Surgery). *J Am Coll Cardiol.* 2007;50:e159-e241.
27. McFalls EO, Ward HB, Moritz TE, et al. Coronary-artery revascularization before elective major vascular surgery. *N Engl J Med.* 2004;351:2795-2804.
28. Poldermans D, Schouten O, Vidakovic R, et al. A clinical randomized trial to evaluate the safety of a noninvasive approach in high-risk patients undergoing major vascular surgery. The DECREASE-V pilot study. *J Am Coll Cardiol.* 2007;49:1763-1769.
29. Harries DJ, Eastwood H. Proximal femoral fractures in the elderly: does operative delay for medical reasons affect short-term outcome? *Age Ageing.* 1991;20:41-44.
30. Sexson SB, Lehner JT. Factors affecting hip fracture mortality. *J Orthop Trauma.* 1987;1:298-305.
31. Geerts WH, Bergqvist D, Pineo GF, et al. Prevention of venous thromboembolism. American college of chest physicians evidence-based clinical practice guidelines (8th edition). *Chest.* 2008;133:381S-453S.
32. Powers PJ, Gent M, Jay RM, et al. A randomized trial of less intense postoperative warfarin or aspirin therapy in the prevention of venous thromboembolism after surgery for fractured hip. *Arch Intern Med.* 1989;149:771-774.
33. Agnelli G, Cosmi B, Di Filippo P, et al. A randomised, double-blind, placebo-controlled trial of dermatan sulphate for prevention of deep vein thrombosis in hip fracture. *Thromb Haemost.* 1992;67:203-208.

34. Lowe GD, Campbell AF, Meek DR, et al. Subcutaneous ancrod in prevention of deep-vein thrombosis after operation for fractured neck of femur. *Lancet*. 1978;2:698-700.
35. Rogers PH, Walsh PN, Marder VJ, et al. Controlled trial of low-dose heparin and sulfinpyrazone to prevent venous thromboembolism after operation on the hip. *J Bone Joint Surg Am*. 1978;60:758-762.
36. Todd CJ, Freeman CJ, Camilleri-Ferrante C, et al. Differences in mortality after fracture of the hip: the east Anglian audit. *BMJ*. 1995;310:904-908.
37. Fisher CG, Blachut PA, Salvian AJ, et al. Effectiveness of pneumatic leg compression devices for the prevention of thromboembolic disease in orthopaedic trauma patients: a prospective, randomized study of compression alone versus no prophylaxis. *J Orthop Trauma*. 1995;9:1-7.
38. Guyatt GH, Cook DJ, Jaeschke R, et al. Grades of recommendation for antithrombotic agents: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines (8th Edition). *Chest*. 2008;133:123S-131S.
39. Borgstom S, Greitz T, van der Linden W, et al. Anticoagulant prophylaxis of venous thrombosis in patients with fractured neck of the femur; a controlled clinical trial using venous phlebography. *Acta Chir Scand*. 1965;129:500-508.
40. Hamilton HW, Crawford JS, Gardiner JH, et al. Venous thrombosis in patients with fracture of the upper end of the femur: a phlebographic study of the effect of prophylactic anticoagulation. *J Bone Joint Surg Br*. 1970;52:268-289.
41. Monreal M, Lafoz E, Navarro A, et al. A prospective double-blind trial of a low molecular weight heparin once daily compared with conventional low-dose heparin three times daily to prevent pulmonary embolus and venous thrombosis in patients with hip fracture. *J Trauma*. 1989;29:873-875.
42. Jorgensen PS, Knudsen JB, Broeng L, et al. The thromboprophylactic effect of a low-molecular-weight heparin (Fragmin) in hip fracture surgery. A placebo-controlled study. *Clin Orthop Relat Res*. 1992;278:95-100.
43. Rosencher N, Vielpeau C, Emmerich J, et al. Venous thromboembolism and mortality after hip fracture surgery: the ESCORTE study. *J Thromb Haemost*. 2005;3:2006-2014.

44. Handoll HH, Farrar MJ, McBirnie J, et al. Heparin, low molecular weight heparin and physical methods for preventing deep vein thrombosis and pulmonary embolism following surgery for hip fractures. *Cochrane Database Syst Rev.* 2002;(4):CD000305.
45. Eriksson BI, Bauer KA, Lassen MR, Turpie AG. Fondaparinux compared with enoxaparin for the prevention of venous thromboembolism after hip-fracture surgery. *N Engl J Med.* 2001;345:1298-1304.
46. Grant PJ, Jaffer AK. When should prophylactic anticoagulation begin after a hip fracture? *Cleve Clin J Med.* 2006;73:785-792.
47. Hefley WF Jr, Nelson CL, Puskarich-May CL. Effect of delayed admission to the hospital on the preoperative prevalence of deep-vein thrombosis associated with fractures about the hip. *J Bone Joint Surg Am.* 1996;78:581-583.
48. Zahn HR, Skinner JA, Porteous MJ. The preoperative prevalence of deep vein thrombosis in patients with femoral neck fractures and delayed operation. *Injury.* 1999;30:605-607.
49. Eriksson BI, Lassen MR; PENTasaccharide in HIp-FRActure Surgery Plus Investigators. Duration of prophylaxis against venous thromboembolism with fondaparinux after hip fracture surgery: a multicenter, randomized, placebo-controlled, double-blind study. *Arch Intern Med.* 2003;163:1337-1342.
50. White RH, Romano PS, Zhou H, et al. Incidence and time course of thromboembolic outcomes following total hip or knee arthroplasty. *Arch Intern Med.* 1998;158:1525-1531.
51. Lawrence VA, Hilsenbeck SG, Noveck H, et al. Medical complications and outcomes after hip fracture repair. *Arch Intern Med.* 2002;162:2053-2057.
52. Zuckerman JD, Skovron ML, Koval KJ, et al. Postoperative complications and mortality associated with operative delay in older patients who have a fracture of the hip. *J Bone Joint Surg Am.* 1995;77:1551-1556.
53. Galanakis P, Bickel H, Gradinger R, et al. Acute confusional state in the elderly following hip surgery: incidence, risk factors and complications. *Int J Geriatr Psychiatry.* 2001;16:349-355.

54. Berggren D, Gustafson Y, Eriksson B, et al. Postoperative confusion after anesthesia in elderly patients with femoral neck fractures. *Anesth Analg*. 1987;66:497-504.
55. Brauer C, Morrison RA, Silberzweig SB, Siu AL. The cause of delirium in patients with hip fracture. *Arch Intern Med*. 2000;160:1556-1860.
56. Gustafson Y, Berggren D, Brannstrom B, et al. Acute confusional states in elderly patients treated for femoral neck fracture. *J Am Geriatr Soc*. 1988;36:525-530.
57. Kalisvaart KJ, de Jonghe JF, Bogaards MJ, et al. Haloperidol prophylaxis for elderly hip-surgery patients at risk for delirium: a randomized placebo-controlled study. *J Am Geriatr Soc*. 2005;53:1658-1666.
58. Marcantonio ER, Flacker JM, Wright RJ, Resnick NM. Reducing delirium after hip fracture: a randomized trial. *J Am Geriatr Soc*. 2001;49:516-522.
59. Johnstone DJ, Morgan NH, Wilkinson MC, Chissell HR. Urinary tract infection and hip fracture. *Injury*. 1995;26:89-91.
60. Michelson JD, Lotke PA, Steinberg ME. Urinary-bladder management after total joint-replacement surgery. *N Engl J Med*. 1988;319:321-326.
61. Skelly JM, Guyatt GH, Kalbfleisch R, et al. Management of urinary retention after surgical repair of hip fracture. *CMAJ*. 1992;146:1185-1189.
62. Beaupre LA, Jones CA, Saunders LD, et al. Best practices for elderly hip fracture patients. A systematic overview of the evidence. *J Gen Intern Med*. 2005;20:1019-1025.
63. Baumgarten M, Margolis D, Berlin JA, et al. Risk factors for pressure ulcers among elderly hip fracture patients. *Wound Rep Reg*. 2003;11:96-103.
64. Hofman A, Geelkerken RH, Wille J, et al. Pressure sores and pressure-decreasing mattresses: controlled clinical trial. *Lancet*. 1994;343:568-571.
65. Siu AL, Penrod JD, Boockvar KS, et al. Early ambulation after hip fracture; effects on function and mortality. *Arch Intern Med*. 2006;166:766-771.
66. National Osteoporosis Foundation. Physician's guide to prevention and treatment of osteoporosis. Washington, DC: National Osteoporosis Foundation; 2003:37.
67. Tang BM, Eslick GD, Nowson C, et al. Use of calcium or calcium in combination with vitamin D supplementation to prevent fractures and bone loss in people aged 50 years and older: a meta-analysis. *Lancet*. 2007;370:657-666.

68. Bischoff-Ferrari HA, Willett WC, Wong JB, et al. Fracture prevention with vitamin D supplementation: a meta-analysis of randomized controlled trials. *JAMA*. 2005;293:2257-2264.
69. AACE Osteoporosis Task Force. American Association of Clinical Endocrinologists medical guidelines for clinical practice for the prevention and treatment of postmenopausal osteoporosis: 2001 edition, with selected updates for 2003. *Endocr Pract*. 2003;9:544-564.
70. Lyles KW, Colon-Emeric CS, Magaziner JS, et al. Zoledronic acid and clinical fractures and mortality after hip fractures. *N Engl J Med*. 2007;357:1799-1809.