

REFERENCES

1. Clement S, Baithwaite SS, Magee MF, et al. Management of diabetes and hyperglycemia in hospitals. *Diabetes Care*. 2004;27:553-591.
2. Van den Berghe G, Wouters P, Weekers F, et al. Intensive insulin therapy in critically ill patients. *N Engl J Med*. 2001;345:1359-1367.
3. Van den Berghe G, Wilmer A, Hermans G, et al. Intensive insulin therapy in the medical ICU. *N Engl J Med*. 2006;354:449-461.
4. Brunkhorst FM, Engel C, Bloos F, et al. Intensive insulin therapy and pentastarch resuscitation in severe sepsis. *N Engl J Med*. 2008;358:125-139.
5. Wiener RS, Weiner DC, Larson RJ. Benefits and risks of tight glucose control in critically ill adults: a meta-analysis. *JAMA*. 2008;300:933-944.
6. The NICE-SUGAR Study Investigators, Finfer S, Chittock DR, et al. Intensive versus conventional glucose control in critically ill patients. *N Engl J Med*. 2009;360:1283-1297.
7. Inzucchi SE, Siegel MD. Glucose control in the ICU-how tight is too tight? *N Engl J Med*. 2009;360:1346-1349.
8. Griesdale DEG, de Souza RJ, van Dam RM, et al. Intensive insulin therapy and mortality among critically ill patients: a meta-analysis including NICE-SUGAR study data. *CMAJ*. 2009;180:821-827.
9. Umpierrez GE, Isaacs SD, Bazargan N, et al. Hyperglycemia: an independent marker of in-hospital mortality in patients with undiagnosed diabetes. *J Clin Endocrinol Metab*. 2002;87:978-982.
10. Furnary A, Zerr K, Grunkemeier G, Starr A. Continuous intravenous insulin infusion reduces the incidence of deep sternal wound infection in diabetic patients after cardiac surgical procedures. *Ann Thorac Surg*. 1999;67:352-362.
11. Furnary AP, Gao G, Grunkemeier GL, et al. Continuous insulin infusion reduces mortality in patients diabetes undergoing coronary artery bypass grafting. *J Thorac Cardiovasc Surg*. 2003;125:1007-1021.
12. Pomposelli JJ, Baxter JK 3rd, Babineau TJ, et al. Early postoperative glucose control predicts nosocomial infection rate in diabetic patients. *J Parenter Enter Nutr*. 1998;22:77-81.
13. McAlister FA, Majumdar SR, Blitz S, et al. The relation between hyperglycemia and outcomes in 2471 patients admitted to the hospital with community-acquired pneumonia. *Diabetes Care*. 2005;28:810-815.
14. Moghissi ES, Korytkowsk MT, DiNardo M, et al. American Association of Clinical Endocrinologists and American Diabetes Association consensus statement on inpatient glycemic control. *Diabetes Care*. 2009;32:1119-1131.
15. Wesorick D, O'Malley C, Rushakoff R, et al. Management of diabetes and hyperglycemia in the hospital: a practical guide to subcutaneous insulin use in the non-critically ill, adult patient. *J Hosp Med*. 2008;3(suppl 5):s17-s28.
16. Inzucchi SE. Management of hyperglycemia in the hospital setting. *N Engl J Med*. 2006;355:1903-1911.
17. Umpierrez GE, Andres P, Smiley D, et al. Randomized study of basal-bolus insulin therapy in the inpatient management of patients with type 2 diabetes (Rabbit 2 trial). *Diabetes Care*. 2007;30:2181-2186.
18. Schnipper JL, Ndumele CD, Liang CL, Pendergrass ML. Effects of a subcutaneous insulin protocol, clinical education, and computerized order set on the quality of inpatient management of hyperglycemia: results of a clinical trial. *J Hosp Med*. 2009;4:16-27.
19. Maynard G, Lee J, Phillips G, et al. Improved inpatient use of basal insulin, reduced hypoglycemia, and improved glycemic control: effect of structured subcutaneous insulin orders and an insulin management algorithm. *J Hosp Med*. 2009;4:3-15.
20. Wesorick DH, Grunawalt J, Gianchandani R. Improving glycemic control in non-critically ill hospitalized patients [abstract]. *J Hosp Med*. 2009;4(suppl 1):63-64.