

REFERENCES

1. Geerts WH, Pineo GF, Heit JA, et al. Prevention of venous thromboembolism: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest*. 2004;126:338S-400S.
2. Shojania KG, Duncan BW, McDonald KM, Wachter RM. Making health care safer: a critical analysis of patient safety practices. Evidence Report/Technology Assessment No. 43. Available at: www.ahrq.gov/clinic/ptsafety/. Accessed July 17, 2007.
3. Practice parameters for the prevention of venous thromboembolism. The Standards Task Force of the American Society of Colon and Rectal Surgeons. *Dis Colon Rectum*. 2000;43:1037-1047.
4. Lindblad B, Eriksson A, Bergqvist D. Autopsy-verified pulmonary embolism in a surgical department: analysis of the period from 1951 to 1988. *Br J Surg*. 1991;78:849-852.
5. Sandler DA, Martin JF. Autopsy proven pulmonary embolism in hospital patients: are we detecting enough deep vein thrombosis? *J R Soc Med*. 1989;82:203-205.
6. Uboch. Statistical Abstract of the United States: 2000. 120th ed. 2000.
7. Rosendaal FR. Venous thrombosis: a multicausal disease. *Lancet*. 1999;353:1167-1173.
8. Stein PD, Henry JW. Prevalence of acute pulmonary embolism among patients in a general hospital and at autopsy. *Chest*. 1995;108:978-981.
9. Agnelli G, Bolis G, Capussotti L, et al. A clinical outcome-based prospective study on venous thromboembolism after cancer surgery: the @RISTOS project. *Ann Surg*. 2006;243:89-95.
10. Mismetti P, Laporte S, Darmon JY, et al. Meta-analysis of low molecular weight heparin in the prevention of venous thromboembolism in general surgery. *Br J Surg*. 2001;88:913-930.
11. Michota F. Controversies and new directions in the management of venous thromboembolism in the perioperative setting. *The Hospitalist*. 2004;6:S32-S38.
12. Perez JV, Warwick DJ, Case CP, Bannister GC. Death after proximal femoral fracture-an autopsy study. *Injury*. 1995;26:237-240.
13. Shojania KG, Duncan BW, McDonald KM, et al. Making health care safer: a critical analysis of patient safety practices. *Evid Rep Technol Assess (Summ)*. 2001;i-x, 1-668.
14. Guyatt G, Schunemann HJ, Cook D, et al. Applying the grades of recommendation for antithrombotic and thrombolytic therapy: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest*. 2004;126:179S-187S.
15. Bergqvist D, Agnelli G, Cohen AT, et al. Duration of prophylaxis against venous thromboembolism with enoxaparin after surgery for cancer. *N Engl J Med*. 2002;346:975-980.
16. Rasmussen MS, Jorgensen LN, Wille-Jorgensen P, et al. Prolonged prophylaxis with dalteparin to prevent late thromboembolic complications in patients undergoing major abdominal surgery: a multicenter randomized open-label study. *J Thromb Haemost*. 2006;4:2384-2390.
17. Amarigiri SV, Lees TA. Elastic compression stockings for prevention of deep vein thrombosis. *Cochrane Database Syst Rev*. 2000:CD001484.
18. Handoll HH, Farrar MJ, McBirnie J, et al. Heparin, low molecular weight heparin and physical methods for preventing deep vein thrombosis and pulmonary embolism following surgery for hip fractures. *Cochrane Database Syst Rev*. 2002:CD000305.
19. Stewart D, Zalamea N, Waxman K, et al. A prospective study of nurse and patient education on compliance with sequential compression devices. *Am Surg*. 2006;72:921-923.
20. Zurawska U, Parasuraman S, Goldhaber SZ. Prevention of pulmonary embolism in general surgery patients. *Circulation*. 2007;115:e302-e307.
21. Agnelli G, Bergqvist D, Cohen AT, et al. Randomized clinical trial of postoperative fondaparinux versus perioperative dalteparin for prevention of venous thromboembolism in high-risk abdominal surgery. *Br J Surg*. 2005;92:1212-1220.
22. Agnelli G, Piovella F, Buoncristiani P, et al. Enoxaparin plus compression stockings compared with compression stockings alone in the prevention of venous thromboembolism after elective neurosurgery. *N Engl J Med*. 1998;339:80-85.
23. Horlocker TT. What's a nice patient like you doing with a complication like this? Diagnosis, prognosis and prevention of spinal hematoma. *Can J Anaesth*. 2004;51:527-534.
24. Eikelboom JW, Quinlan DJ, Douketis JD. Extended-duration prophylaxis against venous thromboembolism after total hip or knee replacement: a meta-analysis of the randomised trials. *Lancet*. 2001;358:9-15.
25. Hull RD, Pineo GF, Stein PD, et al. Extended out-of-hospital low-molecular-weight heparin prophylaxis against deep venous thrombosis in patients after elective hip arthroplasty: a systematic review. *Ann Intern Med*. 2001;135:858-869.
26. Samama CM, Vray M, Barre J, et al. Extended venous thromboembolism prophylaxis after total hip replacement: a comparison of low-molecular-weight heparin with oral anticoagulant. *Arch Intern Med*. 2002;162:2191-2196.
27. Blanchard E, Ansell J. Extended anticoagulation therapy for the primary and secondary prevention of venous thromboembolism. *Drugs*. 2005;65:303-311.
28. Douketis JD, Eikelboom JW, Quinlan DJ, et al. Short-duration prophylaxis against venous thromboembolism after total hip or knee replacement: a meta-analysis of prospective studies investigating symptomatic outcomes. *Arch Intern Med*. 2002;162:1465-1471.
29. Eriksson BI, Lassen MR. Duration of prophylaxis against venous thromboembolism with fondaparinux after hip fracture surgery: a multicenter, randomized, placebo-controlled, double-blind study. *Arch Intern Med*. 2003;163:1337-1342.
30. Rocha AT, de Vasconcellos AG, da Luz Neto ER, et al. Risk of venous thromboembolism and efficacy of thromboprophylaxis in hospitalized obese medical patients and in obese patients undergoing bariatric surgery. *Obes Surg*. 2006;16:1645-1655.
31. Bergqvist D. Risk of venous thromboembolism in patients undergoing cancer surgery and options for thromboprophylaxis. *J Surg Oncol*. 2007;95:167-174.
32. DeBernardo RL, Jr, Perkins RB, Littell RD, et al. Low-molecular-weight heparin (dalteparin) in women with gynecologic malignancy. *Obstet Gynecol*. 2005;105:1006-1011.